

Additivity of the Classical Capacity of Entanglement-Breaking Quantum Channels

Peter W. Shor

AT&T Labs, Florham Park, NJ 07922, USA

February 1, 2008

Abstract

We show that for the tensor product of an entanglement-breaking quantum channel with an arbitrary quantum channel, both the minimum entropy of an output of the channel and the Holevo-Schumacher-Westmoreland capacity are additive. In addition, for the tensor product of two arbitrary quantum channels, we give a bound involving entanglement of formation for the amount of subadditivity (for minimum entropy output) or superadditivity (for classical capacity) that can occur.

One of the more important open questions of quantum information theory is the determination of the capacity of a quantum channel for carrying classical information. This question has been only partially resolved. If entanglement between multiple inputs to the channel is not allowed, a formula for the classical capacity of a quantum channel has indeed been discovered [1, 2]. This capacity formula for a quantum channel Ψ is

$$\chi^*(\Psi) = \max_{p_i, \rho_i} H\left(\sum_i \Psi(p_i \rho_i)\right) - \sum_i p_i H(\Psi(\rho_i)) \quad (1)$$

where H is the von Neumann entropy $H(\rho) = -\text{Tr} \rho \log \rho$, and where the maximization is over probability distributions p_i on density matrices ρ_i over the input space of the channel. This maximum can be attained because we need at most d^2 density matrices ρ_i to achieve any attainable value of

$$\chi(\{p_i, \rho_i\}) = H\left(\sum_i \Psi(p_i \rho_i)\right) - \sum_i p_i H(\Psi(\rho_i)) \quad (2)$$

and are thus maximizing over a compact space. The general capacity of a quantum channel Ψ , without feedback or prior entanglement between sender and receiver, but possibly using entangled inputs, is

$$C(\Psi) = \lim_{n \rightarrow \infty} \frac{1}{n} \chi^*(\Psi^{\otimes n}), \quad (3)$$

i.e., the limit for large n of the capacity when we permit the input to be entangled over blocks of n channel uses. This limit can be shown to exist because χ^* satisfies the superadditivity condition

$$\chi^*(\Psi \otimes \Phi) \geq \chi^*(\Psi) + \chi^*(\Phi). \quad (4)$$

It is conjectured that equality holds, i.e., that χ^* is additive, in which case χ^* would give the classical capacity of a quantum channel without feedback. Substantial work has been done

on this conjecture [3, 4], and it has been proven for several special cases. In particular, it has been proven when one of the channels is the identity channel [4, 5], when one of the channels is what A. S. Holevo calls a c-q or q-c channel (these terms will be defined later) [6, 7], and when one of the channels is a unital qubit channel [8].

We will prove additivity for the special case where one of the two channels is entanglement breaking. Entanglement breaking channels are channels which destroy entanglement with other quantum systems. That is, when the input state is entangled between the input space $\mathcal{H}_{\text{in}}$ and another quantum system $\mathcal{H}_{\text{ref}}$, the output of the channel is no longer entangled with the system $\mathcal{H}_{\text{ref}}$. Both c-q and q-c channels are special cases of entanglement breaking channels. A c-q channel is a channel which can be expressed by the composition of a complete von Neumann measurement on the input space followed by an arbitrary completely positive trace-preserving (CPT) map. A q-c channel can be expressed as the composition of a CPT map followed by a complete von Neumann measurement on the output space. Stated more intuitively, for c-q maps, the input can be treated as being classical, and for q-c maps, the output can be taken to be classical. In either case, the von Neumann measurement eliminates any entanglement between the input space and another system, so c-q and q-c maps are both special cases of entanglement breaking channels. In a conversation with the author, Michal Horodecki [9] gave a simple proof that any entanglement breaking channel can be expressed as a q-c-q channel; that is, the composition of a CPT operator followed by a complete von Neumann measurement followed by another CPT operator. (See also [10] for details of this proof.) As a consequence, the action of an entanglement breaking channel Φ on a state ρ can always be written in the following form introduced by Holevo [6]:

$$\Phi(\rho) = \sum_i \text{Tr}(X_i \rho) \theta_i \quad (5)$$

where $\{X_i\}$ form a general POVM and $\{\theta_i\}$ are arbitrary states. For a c-q map, $X_i = |i\rangle\langle i|$ where $|i\rangle$ form an orthonormal basis, and for a q-c map $\theta_i = |i\rangle\langle i|$.

The additivity problem for capacity is closely related to another additivity problem; that of the minimum entropy output of a channel [11]. For the case of entanglement breaking channels, we first found the additivity proof for the minimum entropy output, and then discovered a straightforward way to extend this additivity proof to cover the classical capacity. In this paper, we first give the proof for additivity of minimum entropy output, as this proof contains the important ideas for the capacity proof, but has significantly fewer technicalities.

Theorem 1 *For an arbitrary quantum channel Ψ , and an entanglement breaking channel Φ*

$$\min_{\rho_{AB}} H((\Psi \otimes \Phi)(\rho_{AB})) = \min_{\rho_A} H(\Psi(\rho_A)) + \min_{\rho_B} H(\Phi(\rho_B)). \quad (6)$$

Proof: The left-hand side is clearly at most the right-hand side, as can be seen by choosing $\rho_{AB} = \rho_A \otimes \rho_B$. We would like to show that it is at least the right-hand side. We use the strong subadditivity property of von Neumann entropy [12]. Consider the minimum obtainable value of $H((\Psi \otimes \Phi)(\rho_{AB}))$. Because Φ is entanglement breaking,

$$(I \otimes \Phi)(\rho_{AB}) = \sum_j q_j |a_j\rangle\langle a_j| \otimes |b_j\rangle\langle b_j| \quad (7)$$

for some q_j , $|a_j\rangle \in \mathcal{H}_A$ and $|b_j\rangle \in \mathcal{H}_B$. Now, we apply to the state

$$\sigma_{ABC} = \sum_j q_j \Psi(|a_j\rangle\langle a_j|) \otimes |b_j\rangle\langle b_j| \otimes |j\rangle\langle j| \quad (8)$$

the property of strong subadditivity in the form

$$H(\sigma_{AB}) \geq H(\sigma_{ABC}) - H(\sigma_{BC}) + H(\sigma_B). \quad (9)$$

We have

$$\begin{aligned} \sigma_{AB} &= \sum_j q_j \Psi(|a_j\rangle\langle a_j|) \otimes |b_j\rangle\langle b_j| \\ &= (\Psi \otimes \Phi)(\rho_{AB}), \end{aligned} \quad (10)$$

the quantity for the entropy of which we would like a lower bound. Now, note that

$$\begin{aligned} H(\sigma_{ABC}) - H(\sigma_{BC}) &= H(\sigma_{AC}) - H(\sigma_C) \\ &= \sum_j q_j H(\Psi(|a_j\rangle\langle a_j|)). \end{aligned} \quad (11)$$

The first equality above follows from the facts that, in Eq. (8), the $|j\rangle\langle j|$ form an orthonormal set and the $|b_j\rangle\langle b_j|$ are pure states, so that $H(\sigma_{ABC}) = H(\sigma_{AC})$ and $H(\sigma_{BC}) = H(\sigma_C)$. The second equality follows from the chain rule for entropy, namely

$$H\left(\sum_j q_j \gamma_j \otimes |j\rangle\langle j|\right) - \sum_j H(q_j |j\rangle\langle j|) = \sum_j q_j H(\gamma_j). \quad (12)$$

Now, note that

$$\begin{aligned} \sigma_B &= \sum_j q_j |b_j\rangle\langle b_j| \\ &= \text{Tr}_A(I \otimes \Phi)(\rho_{AB}) \\ &= \Phi(\text{Tr}_A \rho_{AB}) \end{aligned} \quad (13)$$

Putting the above equalities together, we see that

$$H((\Psi \otimes \Phi)(\rho_{AB})) \geq \sum_j q_j H(\Psi(|a_j\rangle\langle a_j|)) + H(\Phi(\text{Tr}_A \rho_{AB})). \quad (14)$$

Since $\sum_j q_j = 1$, the right-hand side is clearly at least the sum of the minimum output entropies of Ψ and of Φ . We have thus shown that the minimum output entropy is additive for the tensor product of two channels if one of the channels is an entanglement breaking channel. $\square$

We now prove the corresponding additivity result for the Holevo-Schumacher-Westmoreland capacity χ^* ; recall

$$\chi^*(\Psi) = \max_{p_i, \rho_i} H\left(\Psi\left(\sum_i p_i \rho_i\right)\right) - \sum_i p_i H\left(\Psi(\rho_i)\right) \quad (15)$$

over probability distributions p_i and density matrices ρ_i .

Theorem 2 For an arbitrary quantum channel Ψ , and an entanglement breaking channel Φ

$$\chi^*(\Psi \otimes \Phi) = \chi^*(\Psi) + \chi^*(\Phi) \quad (16)$$

Proof: The capacity χ^* is composed of two terms. We will be treating these two terms separately. For the second term, additivity is shown in essentially the same way as in the proof of additivity for minimum entropy, and for the first term, additivity follows from the subadditivity of von Neumann entropy.

Again, we assume that we have an arbitrary quantum channel Ψ , and an entanglement breaking channel Φ . We use strong subadditivity. Consider the optimal signal states for $\Psi \otimes \Phi$, i.e., the p_i and ρ_i such that

$$\chi^*(\Psi \otimes \Phi) = H((\Psi \otimes \Phi)(\rho)) - \sum p_i H((\Psi \otimes \Phi)(\rho_i)) \quad (17)$$

where $\rho = \sum p_i \rho_i$. Let us consider the state $(I \otimes \Phi)(\rho_i)$. Because Φ is an entanglement breaking map, this state is separable, and so

$$(I \otimes \Phi)(\rho_i) = \sum_j q_{ij} |a_{ij}\rangle\langle a_{ij}| \otimes |b_{ij}\rangle\langle b_{ij}| \quad (18)$$

for some $q_{ij}, |a_{ij}\rangle, |b_{ij}\rangle$. Now, we apply strong subadditivity to the state

$$\sigma_{ABC} = \sum_j q_{ij} \Psi(|a_{ij}\rangle\langle a_{ij}|) \otimes |b_{ij}\rangle\langle b_{ij}| \otimes |j\rangle\langle j|. \quad (19)$$

To simplify notation, we let the dependence of σ on i be implicit. Again, we apply strong subadditivity in the form

$$H(\sigma_{AB}) \geq H(\sigma_{ABC}) - H(\sigma_{BC}) + H(\sigma_B) \quad (20)$$

As before,

$$H(\sigma_{AB}) = H((\Psi \otimes \Phi)(\rho_i)). \quad (21)$$

We also have that

$$H(\sigma_B) = H(\Phi(\text{Tr}_A \rho_i)) \quad (22)$$

and

$$H(\sigma_{ABC}) - H(\sigma_{BC}) = \sum_j q_{ij} H(\Psi(|a_{ij}\rangle\langle a_{ij}|)). \quad (23)$$

We let $|a_{ij}\rangle\langle a_{ij}| = \tau_{ij}$. Then $\text{Tr}_B \rho_i = \sum_j q_{ij} \tau_{ij}$. Combining the terms, we observe

$$H((\Psi \otimes \Phi)(\rho_i)) \geq \sum_j q_{ij} H(\Psi(\tau_{ij})) + H(\Phi(\text{Tr}_A \rho_i)) \quad (24)$$

Now, let us sum over all the states ρ_i . We obtain

$$\sum_i p_i H((\Psi \otimes \Phi)(\rho_i)) \geq \sum_{i,j} p_i q_{ij} H(\Psi(\tau_{ij})) + \sum_i p_i H(\Phi(\text{Tr}_A \rho_i)). \quad (25)$$

Using subadditivity of von Neumann entropy and the above inequality (25), we get that

$$\begin{aligned} \chi^*(\Psi \otimes \Phi) &= H((\Psi \otimes \Phi)(\rho)) - \sum_i p_i H((\Psi \otimes \Phi)(\rho_i)) \\ &\leq H(\Psi(\text{Tr}_B \rho)) + H(\Phi(\text{Tr}_A \rho)) \\ &\quad - \sum_{i,j} p_i q_{ij} H(\Psi(\tau_{ij})) - \sum_i p_i H(\Phi(\text{Tr}_A \rho_i)). \end{aligned} \quad (26)$$

However, since

$$\sum_{i,j} p_i q_{ij} \tau_{ij} = \sum_i p_i \text{Tr}_B \rho_i = \text{Tr}_B \rho \quad \text{and} \quad \sum_i p_i \text{Tr}_A \rho_i = \text{Tr}_A \rho, \quad (27)$$

we see that

$$\chi^*(\Psi \otimes \Phi) \leq \chi^*(\Psi) + \chi^*(\Phi). \quad (28)$$

As the opposite inequality is easy, we have additivity of χ^* for entanglement breaking channels. $\square$

We finally give a bound on the amount of superadditivity for general channels. For this, we need to define the entanglement of formation of a bipartite state. This is another quantity that is conjectured to be additive, but for which additivity has not been proved. Entanglement of formation for a bipartite state ρ_{AB} is defined

$$E_F(\rho_{AB}) = \min_{\substack{p_i, \rho_i \\ \sum_i p_i \rho_i = \rho_{AB}}} \sum_i p_i H(\text{Tr}_A \rho_i) \quad (29)$$

where the minimization is over probability distributions p_i on rank-one density matrices ρ_i such that $\sum_i p_i \rho_i = \rho_{AB}$. The theorem is

Theorem 3 *Suppose we have two quantum channels, i.e., completely positive trace preserving maps, Ψ and Φ . Then*

$$\begin{aligned} \min_{\rho_{AB}} H((\Psi \otimes \Phi)(\rho_{AB})) &\geq \min_{\rho_A} H(\Psi(\rho_A)) + \min_{\rho_B} H(\Phi(\rho_B)) \\ &\quad - \max_{\rho_{AB}} E_F((I \otimes \Phi)(\rho_{AB})) \end{aligned} \quad (30)$$

and

$$\chi^*(\Psi \otimes \Phi) \leq \chi^*(\Psi) + \chi^*(\Phi) + \max_{\rho_{AB}} E_F((I \otimes \Phi)(\rho_{AB})). \quad (31)$$

Note that the formulation of the theorem is asymmetric with respect to Ψ and Φ . Thus, to bound the amount of sub- or superadditivity, one can use either the entanglement of formation of $(I \otimes \Phi)(\rho_{AB})$ or of $(\Psi \otimes I)(\rho_{AB})$, whichever is smaller.

Proof: We first give the proof of the first part of Theorem 3. Let

$$(I \otimes \Phi)(\rho_{AB}) = \sum_i q_i \nu_i \quad (32)$$

be the decomposition of $(I \otimes \Phi)(\rho_{AB})$ into pure states ν_i that minimizes entanglement of formation, i.e., so that $\sum_j q_j H(\text{Tr}_A \nu_j)$ is minimum. Now, we consider

$$\sigma_{ABC} = \sum_j q_j (\Psi \otimes I)(\nu_j) \otimes |j\rangle\langle j| \quad (33)$$

and apply strong subadditivity to this state. We obtain

$$H(\sigma_{AB}) \geq (H(\sigma_{ABC}) - H(\sigma_C)) + (H(\sigma_B)) - (H(\sigma_{BC}) - H(\sigma_C)). \quad (34)$$

As in (10), we have

$$H(\sigma_{AB}) = H((\Psi \otimes \Phi)(\rho_{AB})). \quad (35)$$

Similar to (13), we get

$$H(\sigma_B) = H\left(\sum_j q_j \text{Tr}_A \nu_j\right) = H(\Phi(\text{Tr}_A \rho_{AB})) \quad (36)$$

Furthermore, the choice of ν_j and the definition of E_F gives

$$H(\sigma_{BC}) - H(\sigma_C) = E_F\left((I \otimes \Phi)(\rho_{AB})\right) \quad (37)$$

Finally application of the entropy chain rule (12) gives

$$H(\sigma_{ABC}) - H(\sigma_C) = \sum_j q_j H\left((\Psi \otimes I)(\nu_j)\right) \quad (38)$$

The expression (36) is bounded below by $\min_\rho H(\Phi(\rho))$. The second expression (37) is bounded above by $\max_\rho E_F((I \otimes \Phi)(\rho))$. The third expression (38) is bounded below by $\min_\rho H((\Psi \otimes I)(\rho))$, which is known to equal $\min_\rho H(\Psi(\rho))$. Combining these three expressions give the first part of Theorem 3.

To prove the second part of the theorem, (38) must be replaced by

$$H(\sigma_{ABC}) - H(\sigma_C) \geq \sum_{jk} q_j r_{jk} H\left(\Psi(|v_{jk}\rangle\langle v_{jk}|)\right) \quad (39)$$

for states $|v_{jk}\rangle$ and probabilities $q_j r_{jk}$ such that

$$\sum_{j,k} q_j r_{jk} |v_{jk}\rangle\langle v_{jk}| = \text{Tr}_B \nu_j. \quad (40)$$

We then consider the signal states ρ_i and the associated probabilities p_i which give the value of $\chi^*(\Psi \otimes \Phi)$ in Equation (1), and let $\sum_i p_i \rho_i = \rho$. We now use expressions (36), (37), (39) with ρ_i in the place of ρ_{AB} . Combining these three expressions yields

$$\begin{aligned} H\left((\Psi \otimes \Phi)(\rho_i)\right) &\geq H\left(\Phi(\text{Tr}_A \rho_i)\right) + \sum_{j,k} q_{ij} r_{ijk} H\left(\Psi(|v_{ijk}\rangle\langle v_{ijk}|)\right) \\ &\quad - E_F\left((I \otimes \Phi)(\rho_i)\right). \end{aligned} \quad (41)$$

The second part of Theorem 3 then follows in a way entirely analogous to the proof of Theorem 2. We use the equalities

$$\text{Tr}_B \rho = \sum_i p_i \text{Tr}_A \rho_i \quad (42)$$

and

$$\text{Tr}_A \rho = \sum_{i,j,k} p_i q_{ij} r_{ijk} |v_{ijk}\rangle\langle v_{ijk}|, \quad (43)$$

and expand $\chi^*(\Psi \otimes \Phi)$ similarly to Eq. (26) to obtain Eq. (31).

We still must prove the inequality (39). The left hand side of (39) is

$$\sum_j q_j H\left((\Psi \otimes I)(\nu_j)\right) \quad (44)$$

Now, ν_j is a purification of $\text{Tr}_B \nu_j$, and $H((\Psi \otimes I)(\nu_j)) = H((\Psi \otimes I)(\tau))$ for any other quantum state τ which is a purification of $\sigma_j = \text{Tr}_B \nu_j$. Let $\sigma_j = \sum_k q_{jk} |v_{jk}\rangle\langle v_{jk}|$ be the eigenvector decomposition of σ_j . A different purification is

$$\tau_j = \left(\sum_k q_{jk} |v_{jk}\rangle \otimes |k\rangle \otimes |k\rangle\right) \left(\sum_k q_{jk} \langle v_{jk}| \otimes \langle k| \otimes \langle k|\right) \quad (45)$$

It suffices to show that

$$H((\Psi \otimes I)(\tau_j)) \geq H(\text{Tr}_3(\Psi \otimes I)(\tau_j)) - H(\text{Tr}_{12}(\Psi \otimes I)(\tau_j)) \quad (46)$$

as the first term in the above equation is $H((\Psi \otimes I)(\nu_j))$, the second is $H(\{q_{jk}\}_k) + \sum_k q_{jk} H(\Psi(|v_{jk}\rangle\langle v_{jk}|))$, and the third is $H(\{q_{jk}\}_k)$. However, the above equation follows from the inequality $H(\rho_{34}) \geq H(\rho_3) - H(\rho_4)$, which is a consequence (after another purification) of the subadditivity property of entropy [13]. $\square$

Acknowledgements

I would like to thank Michal and Pawel Horodecki, Chris King, and Beth Ruskai for interesting and informative conversations. In addition, I would like to thank Michal Horodecki for discovering the proof that all entanglement-breaking channels can be expressed in Holevo's form (i.e., Eq. (5)), and Chris King and Beth Ruskai for their extremely helpful suggestions for improving the exposition of this paper.

References

- [1] A. S. Holevo, "The capacity of quantum channel with general signal states," *IEEE Trans. Info Thy.* **44**, 269–273 (1998); LANL e-print quant-ph/9611023.
- [2] B. Schumacher and M. D. Westmoreland, "Sending classical information via noisy quantum channels," *Phys. Rev. A* **56**, 131–138 (1997).
- [3] G. G. Amosov and A. S. Holevo, "On the multiplicativity conjecture for quantum channels," LANL e-print math-ph/0103015 (2001).
- [4] G. G. Amosov, A. S. Holevo, and R. F. Werner "On some additivity problems in quantum information theory," *Probl. Inform. Transm.* **36** (4), 25–34 (2000); LANL e-print math-ph/0003002.
- [5] B. Schumacher and M. D. Westmoreland, "Relative entropy in quantum information theory," in *Quantum Information and Quantum Computation: A Millenium Volume*, AMS Contemporary Mathematics Vol. 305, (S. J. Lomomaco, Jr. and H. E. Brandt, eds., AMS Press, Providence, Rhode Island, 2002); LANL e-print quant-ph/0004045.
- [6] A. S. Holevo, "Coding theorems for quantum channels," *Russian Math Surveys* **53**, 1295–1331 (1998); LANL e-print quant-ph/9809023.
- [7] C. King, "Maximization of capacity and l_p norms for some product channels," *J. Math. Phys* **43**, 1247–1260 (2002); LANL e-print quant-ph/0103086.
- [8] C. King, "Additivity for unital qubit channels," LANL e-print quant-ph/0103156 (2001).
- [9] M. Horodecki, private communication (2001).
- [10] M. B. Ruskai, "Entanglement breaking channels," *Rev. Math. Phys.*, to appear; LANL e-print quant-ph/0207100.
- [11] C. King and M. B. Ruskai, "Minimal entropy of states emerging from noisy quantum channels," *IEEE Trans. Inf. Theory* **47**, 1–19 (2001); LANL e-print quant-ph/0103156.
- [12] E. Lieb and M.B. Ruskai, "Proof of the strong subadditivity of quantum mechanical entropy," *J. Math. Phys.* **14**, 1938–1941 (1973).
- [13] H. Araki and E. Lieb, "Entropy inequalities," *Commun. Math. Phys.* **18**, 160–170 (1970).